

Targeting Inflammation to Slow CKD Progression- Role of Gut Microbiome Based Therapy

Emmanuel Anteyi*, Natarajan Ranganathan and Usha Vyas

Clinical Research & Development, Kibow Therapeutics, Inc 4781 West Chester Pike, Newtown Square, PA 19073, USA

Abstract

Chronic Kidney Disease (CKD) is a worldwide public health problem associated with high morbidity, mortality, and socioeconomic burden. Despite recent advances in pathogenesis and effective Standard of Care (SOC) therapy, the incidence of end stage renal disease remains high, attributed to the complex and diverse pathophysiological pathways in progression of CKD. Among the many pathogenetic pathways, inflammation plays the most central role in disease progression and related complications. Current SOC treatment to slow CKD progression has not adequately addressed these complex processes as a result of persistence of low-grade inflammation occurring at all stages of CKD. Previous attempts to address the inflammatory pathway with specific anti-inflammatory agents were, in certain cases discontinued for reasons ranging from drug safety to efficacy concerns, and business development decisions. Recent research findings demonstrating direct correlation between systemic inflammation associated gut microbiota changes and CKD progression, provided a potential novel therapeutic approach to target the inflammatory pathways. Exploring current scientific knowledge of modulating gut microbiome with microbiome-based therapies provides options in addressing these complex pathogenetic mechanisms of CKD-induced inflammation.

The purpose of this minireview is to discuss the role of inflammation associated with gut microbiome changes in CKD progression and gut microbiome-based therapeutic options to slow CKD progression.

Keywords: CKD progression • Inflammation • Gut microbiome

Introduction

Chronic Kidney Disease (CKD) is a worldwide public health problem associated with high morbidity, mortality and disproportionately higher healthcare expenditure compared to other disease conditions [1]. CKD has been projected to be among leading top five causes of death worldwide attributed to elderly population and high prevalence of cardiovascular complications from diabetes [2]. CKD progression is characterized by persistent low-grade systemic inflammation typified by presence of proinflammatory markers, similar to what obtains in other chronic inflammatory conditions such as metabolic syndrome, Non-Alcoholic Fatty Liver Disease (NAFLD), Cardiovascular Disease (CVD), malignancy and Diabetes Mellitus (DM) [3]. The inflammatory process is initiated by tissue injury or presence of foreign particles which triggers production of proinflammatory markers culminating in inflammatory cascade. This physiological proinflammatory process is controlled or abated by counter regulatory production of anti-inflammatory molecules and any residual and persistent low-grade inflammation is either due to failure to discontinue the inflammatory cascade or there is a continued generation of ongoing pro-inflammatory markers [3,4].

This persistent inflammatory phenomenon was also found in conditions associated with poor nutritional status, gut microbiota dysbiosis, infections, dyslipidemia, and stress [4]. Inflammation plays significant components in uremic toxemia, oxidative stress, infections, dyslipidemia, malnutrition, volume

overload and dialysis treatment, which are well known contributing factors to CKD progression and related cardiovascular complications [4,5]. Targeting inflammatory pathway plays a key role in therapeutic approach in CKD progression such as Renin-Aldosterone Angiotensin System (RAAS) blockers, Sodium Glucose Co-transporter 2 inhibitor (SGLT2i), Mineralocorticoid Receptor Antagonist (MRA) and other anti-inflammatory agents undergoing clinical development [6]. Despite evidence of clinical benefits of these medications as Standard Of Care (SOC) therapy, risk for disease progression persists, necessitating further search for a safe novel therapy targeting inflammation in high-risk patients for CKD progression.

Recent evidence of systemic inflammation in CKD-associated gut dysbiosis has provided further mechanistic insight into the complex relationship between inflammation and molecular, immunological, and metabolic pathways in CKD progression [7,8]. Exploring gut microbiota modulation targeting inflammatory process to slow CKD progression is a pivotal intervention central to these complex and interrelated pathophysiological phenomena [9]. The purpose of this review is to briefly summarize the key roles of inflammation and gut microbiome changes in CKD progression and discuss gut microbiome-based therapy options to slow CKD progression.

Inflammation and CKD progression

Recent reviews on therapeutic options to retard progression of kidney diseases highlighted the central role of inflammation in initiation and outcome of CKD, including its significant relationship with pathophysiological pathways of hemodynamic, metabolic, and immunologic factors [4,6,10]. Potential therapy to slow CKD progression would, therefore, be multi-approach, preferably combination therapies to target the complex pathways addressing the various incompletely understood pathogenetic mechanisms [10]. The need for novel multi-approach to inflammation of CKD was further supported by recent findings that despite available effective agents such as RAAS blockers, SGLT2 inhibitors and MRA as standard of care therapy for kidney protection, a significant proportion of patients still develop ESRD [6,11,12]. This points to the fact that treatment gaps exist and significant limitations of clinical benefits by current SOC therapy in CKD. The unmet needs in high-risk patients for CKD progression necessitate discovery and development of novel therapies targeting inflammatory pathways.

***Address for Correspondence:** Emmanuel Anteyi, Clinical Research & Development, Kibow Therapeutics, Inc 4781 West Chester Pike, Newtown Square, PA 19073, USA, Tel: 5714300753; E-mail: e.anteyi@kibowbiotech.com

Copyright: © 2024 Anteyi E, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Received: 01 January, 2024; Manuscript No. jnt-23-118437; **Editor Assigned:** 02 January, 2024; PreQC No. P-118437; **Reviewed:** 17 January, 2024; QC No. Q-118437; **Revised:** 23 January, 2024, Manuscript No. R-118437; **Published:** 31 January, 2024, DOI: 10.37421/2161-0959.2024.14.481

Inflammation is a well-recognized pathway in CKD pathogenesis as demonstrated by reduced disease progression in response to intervention with anti-inflammatory agents [13,14]. Despite the potential benefits, clinical development of some of these anti-inflammatory agents were discontinued for reasons ranging from drug safety concerns to uncertain future investment returns and recent approval of SGLT2i and MRA in CKD treatment, raised the required standard for potential pipeline products in treatment of kidney disease progression [6,15]. Recent advances in gut microbiome-based therapy approaches provides additional opportunity to explore safe and effective novel anti-inflammatory to curb the high CKD and ESRD incidence [16,17].

The principle of inflammation is based on detection and elimination of harmful pathogens through interaction of resident renal parenchymal cells and various immune cells, a process initiated through either stress or injury, leading to irreversible tissue damage and organ dysfunction [10,18]. In the kidneys, the initial cellular interactions are between resident parenchymal immune cells (macrophages, dendritic cells) and circulating immune cells (monocytes, lymphocytes, neutrophils) [19]. The exact mechanistic process involved in inflammation of CKD progression is not well understood, but is known to include contribution from hemodynamic, immunologic, and metabolic etiological factors [20].

Further findings showed that even CKD that is not immune mediated in etiology, has inflammation playing a significant pathogenetic role in disease progression [21]. Initiation of inflammatory process depends mainly on resident immune cells whose key responsibility is to maintain tissue homeostasis between the Dendritic Cells (DC), macrophages, regulatory T cells (Tregs), CD8, NK lymphocytes, who are closely in contact and interact with renal parenchymal cells [22]. Once these cells are triggered by kidney injury from external or internal agents (microbial antigens, toxins, hyperglycemia, proteinuria) produce inflammatory mediators, initiating inflammatory cascade that sets the process of kidney disease progression. As a result of initiation of inflammatory process, a counter-regulatory physiological response is triggered to control inflammation, repair tissue damage, and restore homeostasis [23].

This key initiating step involve activation of DC, leading to enhance in activity of CD8+T, CD4+T and TH2 causing glomerulopathy with infiltrations by macrophages thereby amplify repeated processes of repair and fibrotic changes [4,10,23,24]. The onset of glomerulopathy and proteinuria triggers RAAS which further increased production of several proinflammatory factors, cytokines, chemokines, and growth factors (Figure 1). These secondary responses activate innate immunity and signaling transcription factors particularly NF- κ B, NLRP3 inflammasome, TLR, Nrf2.

The activated transcriptomes regulate proinflammatory and senescence factors (IL-6, TNF α , CCR2, CCR5, JAK-STAT2, MCP1, α -klotho), thereby setting a vicious cascade of aggravated inflammation and renal fibrosis [23,25] (Figure 1). Further insight in recent Acute Kidney Injury (AKI) and CKD models showed the inflammatory process in AKI persists and continues to CKD, even after the renal function was restored, confirming strong association between low-grade inflammation and slow development of disease progression and fibrosis [26,27]. Mechanistic studies of how the activation of these cells and receptors and outcomes measures are potential targets for novel therapeutic agents in CKD progression. Clear understanding of inflammatory pathways triggered by uremic toxins forms the basis to explore benefits of Live Biotherapeutic Products (LBP) as gut microbiome-based therapy in CKD progression [28].

Gut microbiome in CKD progression

The human gastrointestinal tract harbors complex commensal microorganisms of bacteria, archaea, small eukaryotes, and viruses called microbiomes. The human microbiome is predominantly located in the colon consisting mainly of diverse bacteria, whose cell composition is 10 times greater than human cells [29]. Factors affecting gut microbiota composition and function include host factors such as mode of birth delivery, gestational age, breastfeeding, age, diet, geographical location, antibiotic use, sanitation [28]. Various studies showed microbiome-host interactions modulate many vital functions in healthy human host such as metabolism, immunity, cardiovascular and neurological functions [30]. The gut microbiota and host




Figure 1. Summary of key roles of inflammation in initiation and progression of chronic kidney disease. Abbreviations: DC: Dendritic Cells; MC: Macrophages; IL: Interleukins; TNF: Tumor Necrosis Factor; GM-CSF: Granulocyte-Macrophage Colony Stimulating Factor; TGF: Transfer Growth Factor; NF- κ B: Nuclear Factor-Kappa B; MCP1: Monocyte Chemoattractant Protein1; CCR2/CCR5: CC-Chemokine Receptor; Nrf2-Keap1: Nuclear Factor-(erythroid -derived 2)- Like 2; Keap 1: Kelch-Like ECH-Associated protein 1; ASK1: Apoptosis Signal-regulating Kinase 1, JAK-STAT2:Janus Kinase/Signal Transducers and Activators of Transcription; PDE: Phosphodiesterase; ECM: Extracellular Matrix; EMT: Epithelial-to-Mesenchymal Transition.

are in a mutualistic relationship with microbiota utilizing host nutrients in a conducive environment and at same time modulating large vital functions of host immunity and metabolic process [31]. Additionally, these bacteria encode enzymes to produce metabolites essential to host health, such as vitamins, bile acids, choline, and Short Chain Fatty Acids (SCFA).

The SCFA are fermentation products of non-digestible dietary carbohydrate by gut microbiota consisting mainly of butyrate, propionate, and acetate whose main functions are provision of energy source for colonocytes, maintain integrity of intestinal barrier, regulates glucose and lipid metabolism, and regulates immune system [32]. Specific aspects of immune regulation involve attenuation of NF- κ B activation and inhibition of proinflammatory cytokine production through activity of Tregs cells, protecting against sustained activation of immune system [33,34]. In addition, SCFA inhibits histone deacetylases through their action on G-Protein Coupled Receptors which includes GPR41 (FFAR3), GPR43 (FFAR2) and GPR109a responsible for regulating various functions and metabolic effects of adipocytes, immune and vascular endothelial cells [35]. Persistent and low-grade inflammation is a known process associated with many chronic diseases like CKD, cardiovascular, metabolic syndrome, neurological and allergic disorder [36].

The initiation of inflammation starts with gut microbial-host interaction, resulting in functional and compositional changes of the four main microbiota phyla, namely Actinobacteria, Proteobacteria, Firmicutes and Bacteroidetes

[37]. Evidence indicates that changes in microbiota composition and function (gut dysbiosis) causes gut inflammation and disruption of intestinal barrier integrity, ("leaky bowel") results in activation of NF- κ B pathway and eventual development of systemic inflammation [38]. Consequences of disrupted intestinal barrier includes translocation of bacterial toxins, Lipopolysaccharide (LPS), uremic toxins and endotoxemia with increased production of proinflammatory cytokines and gut-derived uremic toxins (IS, PCS, TMAO) which further worsens inflammation, increases proinflammatory cytokines and Reactive Oxygen Species (ROS) production [39,40]. High levels of uremic toxins have been implicated in pathogenesis of renal fibrosis and cell senescence through mechanistic reduction of klotho expression [41,42].

The relationship between kidney dysfunction and gut dysbiosis is bidirectional, termed "Gut-kidney axis" in which CKD causes gut microbial changes (dysbiosis) and in turn worsens CKD, aggravated by complications of metabolic acidosis, hypervolemia, prolonged colonic transit time, antibiotic use, intestinal wall edema and poor dietary fiber intake [10,43-45]. The systemic inflammation induced by gut dysbiosis is characterized by increased composition of colon bacteria families possessing enzymes that generate uremic toxins compared to dietary fiber degraders that produce beneficial SCFA metabolite, given rise to reduced SCFA production, accumulation of harmful metabolites and endotoxemia prior to systemic inflammation [46,47] (Figure 2).

Targeting inflammation with gut microbiome-based therapy

Previous interventions in CKD progression directly targeting various specific inflammatory markers have not significantly reduced high CKD burden [48]. Various mechanistic studies have shown that current recommended RAAS blockers, SGLT2i, MRA and GLP-1 receptor agonist anti-inflammatory benefits were attributed to secondary reduction of glomerular hyperfiltration and proteinuria [49-51]. Additionally, there are ongoing clinical development of various therapeutic agents targeting specific inflammatory markers such as Nrf2 activators, Endothelin-1 Receptor Agonist (ERA), Soluble Guanyl Cyclase Activator(sGC), Anti-inflammatory cytokines (anti-TNF, IL-6, CCL2, NF- κ B) and cell therapy [52-54].

Several studies have reported that CKD -associated gut microbiome

changes have a direct relationship with disease progression and live biotherapeutic interventions to restore colonic bacteria balance (eubiosis) are known to improve biochemical and clinical outcomes in CKD [55,56]. Targeting CKD- induced inflammation with microbiome-based therapy addresses pathways that reduce uremic toxins, increase beneficial SCFA metabolites, restores epithelial barrier, regulates immune system, and decreases persistent low-grade inflammation [57].

The dietary approach is well known cost-effective approach to modulate gut microbiome with recommendation that support plant-based diets and low animal protein to decrease bacterial proteolytic fermentation while promoting and increasing saccharolytic bacterial composition. This dietary regime affects microbiota composition that reduces inflammation and uremic toxin production associated with CKD progression and CVD complications [58,59]. The goal of gut-based microbiome intervention primarily is normalization of CKD-associated dysbiosis by reducing composition and activity of proteolytic bacterial species while increasing that of saccharolytic species, classified as prebiotics, probiotics and synbiotics [60]. Prebiotics are non-digestible carbohydrate fibers selectively used as substrates by host microorganisms to confer a health benefit (examples of dietary fiber are inulin, resistant starch, fructo- and galacto-oligosaccharide) [61].

Probiotics are live microorganisms, which when administered in adequate amounts confer a health benefit on the host (examples of probiotic strains include, Bifidobacterium, Lactobacillus and Streptococcus species) [62]. Several studies reported probiotics significantly reduce gut derived uremic toxins, inflammatory markers and proinflammatory cytokines in all CKD stages. These clinical benefits corresponded to changes in colonic bacteria taxa providing further proof of gut-kidney involvement in CKD progression. Synbiotics is a combined therapy of prebiotics and probiotics synergistically reduce the effect of uremic toxins and ameliorate gut dysbiosis in CKD progression [63]. Intervention studies in advanced CKD using combination prebiotic (Inulin, fructo-oligosaccharide, galacto-oligosaccharide) and probiotic (lactobacilli, streptococci, Bifidobacterium) showed altered composition of fecal microbiome with increased Bifidobacterium, and while pathogenic Ruminococaceae were reduced [64,65]. In addition, various synbiotic interventions were reported to decrease protein bound uremic toxins such as




Figure 2. Graphic illustration demonstrating complex relationship between CKD-associated gut microbiome changes (Dysbiosis) and Inflammation in development of chronic kidney disease progression. Adapted with Permission from: Mertowska P, Mertowski S, Wojnicka J, et al (2021). Link between Chronic Kidney Disease and Gut Microbiota in Immunological and Nutritional Aspects. *Nutrients.*; 13(10):3637. <https://doi.org/10.3390/nu13103637>

p-cresol in Pre-dialysis and dialysis CKD populations [66].

A new concept of postbiotic as adjunctive or alternative therapeutic approach in gut homeostasis and immune modulation was recently proposed and the term postbiotic refers to products of non-viable bacteria or metabolites of probiotic organisms such as vitamins, SCFA, cell surface proteins and enzymes that have demonstrated positive effect on gut microbiome and host [67]. In contrast to synbiotic products, postbiotics have pharmacokinetic properties of absorption, distribution, metabolism, and excretion raising concerns on dosage and systemic toxic effects. An experimental postbiotic SCFA treatment demonstrated reduction of inflammation and oxidative stress in ischemia-reperfusion animal model of kidney injury [68].

Another Live Biotherapeutic Product (LBP) approach is fecal microbiota transplantation, as capsules or through colonoscopy, normally sourced from healthy donors with preclinical data showing potential benefits in CKD patients. This mode of therapy has recently gotten approval for recurrent *Clostridium difficile* infection and clinical benefits have equally been reported in metabolic syndrome, autism, IBS, and ESRD-associated bacteremia [69].

In comparison to drug therapy, some of the microbiome-based therapy is cheaper with well tolerated fewer side effects across spectrum of CKD/ESRD patients.

Conclusion

Inflammatory process plays a central role in the initiation and sustenance of CKD progression. The persistence of low-grade systemic inflammation in CKD is partly a reason partly given for the high burden of CKD and related CVD complications. Despite decades of available effective standard of care therapy to slow CKD progression such as RAAS blockers, SGLT2 inhibitor, MRA and GLP-1 agonist, the high incidence of CKD/ESRD still persists. The inability to reduce CKD public health burden showed that current interventions are inadequate to address the complex and interrelated pathogenetic pathways, particularly the key and central role of inflammation in initiation and progression of kidney disease. Hence, further clinical studies to identify novel therapies targeting inflammation will bridge scientific gap and address unmet needs of patients who are high risk for CKD progression. Gut dysbiosis is associated with systemic inflammation in CKD and hence microbiome modulation through microbiota-based therapy is a potential anti-inflammatory therapeutic product in retarding disease progression.

Acknowledgement

Sumanth Narayana helped with graphical and digital illustrations of Figures 1 and 2.

Disclosures

No conflict of interest.

Authors Contribution

Conceptualization: Natarajan Ranganathan

Writing-original draft: Emmanuel Anteyi

Writing-review & editing: Emmanuel Anteyi, Usha Vyas

Data Sources

NA.

Funding

None.

IRB Approval

NA.

References

- Kovesdy, Csaba P. "Epidemiology of chronic kidney disease: An update 2022." *Kidney Int Suppl* 12 (2022): 7-11.
- Hill, Nathan R., Samuel T. Fatoba, Jason L. Oke and Jennifer A. Hirst, et al. "Global prevalence of chronic kidney disease-A systematic review and meta-analysis." *PLoS one* 11 (2016): e0158765.
- United States Renal Data System. 2022 USRDS Annual Data Report: Epidemiology of kidney disease in the United States. National institutes of health; National institute of diabetes and digestive and kidney diseases; Bethesda, MD, USA.
- Saurav, Prashant Kadam, Matthew Satariano, Michael Massey and Kai Mongan, et al. "The role of inflammation in CKD." *Cells* 12 (2023): 1581.
- Mihai, Simona, Elena Codrici, Ionela Daniela Popescu and Ana-Maria Enciu, et al. "Inflammation-related mechanisms in chronic kidney disease prediction, progression and outcome." *J Immunol Res* 2018 (2018).
- Rayego-Mateos, Sandra, Raul Rodriguez-Diez, Beatriz Fernandez-Fernandez and Carmen Mora-Fernández, et al. "Targeting inflammation to treat diabetic kidney disease: The road to 2030." *Kidney Int* (2022).
- Lau, Wei Ling, Kamyar Kalantar-Zadeh and Nosratola D. Vaziri. "The gut as a source of inflammation in chronic kidney disease." *Nephron* 130 (2015): 92-98.
- Lau, Wei Ling, Javad Savoj, Michael B. Nakata and Nosratola D. Vaziri. "Altered microbiome in chronic kidney disease: Systemic effects of gut-derived uremic toxins." *Clin Sci* 132 (2018): 509-522.
- Nallu, Anitha, Shailendra Sharma, Ali Ramezani and Jagadeesan Muralidharan, et al. "Gut microbiome in chronic kidney disease: Challenges and opportunities." *Transl Res* 179 (2017): 24-37.
- Andrade-Oliveira, Vinicius, Orestes Foresto-Neto, Ingrid Kazuo Mizuno Watanabe and Roberto Zatz, et al. "Inflammation in renal diseases: New and old players." *Front pharmacol* 10 (2019): 1192.
- American Diabetes Association Professional Practice Committee. "Chronic kidney disease and risk management: Standards of medical care in diabetes-2022." *Diabetes care* 45 (2022): S175-S184.
- JJ, Garcia Sanchez, J. Thompson, D. A. Scott and R. Evans, et al. "Treatments for chronic kidney disease: A systematic literature review of randomized controlled trials." *Adv Ther* 39 (2021): 193-220.
- Akchurin, Oleh M and Frederick Kaskel. "Update on inflammation in chronic kidney disease." *Blood Purif* 39 (2015): 84-92.
- Machowska, Anna, Juan Jesus Carrero, Bengt Lindholm and Peter Stenvinkel. "Therapeutics targeting persistent inflammation in chronic kidney disease." *Transl Res* 167 (2016): 204-213.
- Hofherr, Alexis, Julie Williams, Li-Ming Gan and Magnus Söderberg, et al. "Targeting inflammation for the treatment of diabetic kidney disease: A five-compartment mechanistic model." *BMC Nephrol* 23 (2022): 208.
- Ranganathan, Natarajan and Emmanuel Anteyi. "The role of dietary fiber and gut microbiome modulation in progression of chronic kidney disease." *Toxins* 14 (2022): 183.
- Hobby, Gerren P., Oleg Karaduta, Giuseppina F. Dusio and Manisha Singh, et al. "Chronic kidney disease and the gut microbiome." *Am J Physiol Renal Physiol* 316 (2019): F1211-F1217.
- Singbartl, Kai, Cassandra L. Formeck and John A. Kellum. "Kidney-immune system crosstalk in AKI." *Semin Nephrol* 39 (2019) 96-106.
- Kurts, Christian, Ulf Panzer, Hans-Joachim Anders and Andrew J. Rees. "The immune system and kidney disease: Basic concepts and clinical implications." *Nat Rev Immunol* 13 (2013): 738-753.
- Zoccali, Carmine, Raymond Vanholder, Ziad A. Massy and Alberto Ortiz, et al. "The systemic nature of CKD." *Nat Rev Nephrol* 13 (2017): 344-358.

21. Floege, Jürgen, Mark W. Burns, Charles E. Alpers and Ashio Yoshimura, et al. "Glomerular cell proliferation and PDGF expression precede glomerulosclerosis in the remnant kidney model." *Kidney Int* 41 (1992): 297-309.
22. Heymann, Felix, Catherine Meyer-Schwesinger, Emma E. Hamilton-Williams and Linda Hammerich, et al. "Kidney dendritic cell activation is required for progression of renal disease in a mouse model of glomerular injury." *J Clin Invest* 119 (2009): 1286-1297.
23. Stenvinkel, Peter, Glenn M. Chertow, Prasad Devarajan and Adeera Levin, et al. "Chronic inflammation in chronic kidney disease progression: Role of Nrf2." *Kidney Int Rep* 6 (2021): 1775-1787.
24. Lv, Wenshan, George W. Booz, Yangang Wang and Fan Fan, et al. "Inflammation and renal fibrosis: Recent developments on key signaling molecules as potential therapeutic targets." *Eur J Pharmacol* 820 (2018): 65-76.
25. Turner, Clare M., Nishkantha Arulkumaran, Mervyn Singer and Robert J. Unwin, et al. "Is the inflammasome a potential therapeutic target in renal disease?" *BMC Nephrol* 15 (2014): 1-13.
26. Chawla, Lakhmir S., Paul W. Eggers, Robert A. Star and Paul L. Kimmel. "Acute kidney injury and chronic kidney disease as interconnected syndromes." *N Engl J Med* 371 (2014): 58-66.
27. Zuk, Anna and Joseph V. Bonventre. "Recent advances on acute kidney injury and its consequences and impact on chronic kidney disease." *Curr Opin Nephrol Hypertens* 28 (2019): 397.
28. Krukowski, Hubert, Sophie Valkenburg, Avra-Melina Madella and Johan Garssen, et al. "Gut microbiome studies in CKD: Opportunities, pitfalls and therapeutic potential." *Nat Rev Nephrol* 19 (2023): 87-101.
29. Kho, Zhi Y and Sunil K. Lal. "The human gut microbiome—a potential controller of wellness and disease." *Front Microbiol* 9 (2018): 1835.
30. Fan, Yong and Oluf Pedersen. "Gut microbiota in human metabolic health and disease." *Nat Rev Microbiol* 19 (2021): 55-71.
31. Vanholder, Raymond and Griet Glorieux. "The intestine and the kidneys: A bad marriage can be hazardous." *Clin Kidney J* 8 (2015): 168-179.
32. Rowland, Ian, Glenn Gibson, Almut Heinken and Karen Scott, et al. "Gut microbiota functions: Metabolism of nutrients and other food components." *Eur J Nutr* 57 (2018): 1-24.
33. Chang, Pamela V., Liming Hao, Stefan Offermanns and Ruslan Medzhitov. "The microbial metabolite butyrate regulates intestinal macrophage function via histone deacetylase inhibition." *Proc Natl Acad Sci* 111 (2014): 2247-2252.
34. Smith, Patrick M., Michael R. Howitt, Nicolai Panikov and Monia Michaud, et al. "The microbial metabolites, short-chain fatty acids, regulate colonic T_{reg} cell homeostasis." *Science* 341 (2013): 569-573.
35. Hara, Takafumi, Ikuro Kimura, Daisuke Inoue and Atsuhiko Ichimura, et al. "Free fatty acid receptors and their role in regulation of energy metabolism." *Rev Physiol Biochem Pharmacol* 164 (2013): 77-116.
36. Chen, Yinwei, Jinghua Zhou and Li Wang. "Role and mechanism of gut microbiota in human disease." *Front Cell Infect Microbiol* 11 (2021): 86.
37. Carding, Simon, Kristin Verbeke, Daniel T. Vipond and Bernard M. Corfe, et al. "Dysbiosis of the gut microbiota in disease." *Microb Ecol Health Dis* 26 (2015): 26191.
38. Anders, Hans-Joachim, Kirstin Andersen and Bärbel Stecher. "The intestinal microbiota, a leaky gut, and abnormal immunity in kidney disease." *Kidney Int* 83 (2013): 1010-1016.
39. Rysz, Jacek, Beata Franczyk, Janusz Ławiński and Robert Olszewski, et al. "The impact of CKD on uremic toxins and gut microbiota." *Toxins* 13 (2021): 252.
40. Graboski, Amanda L and Matthew R. Redinbo. "Gut-derived protein-bound uremic toxins." *Toxins* 12 (2020): 590.
41. Yang, Yi, Milos Mihajlovic and Rosalinde Masereeuw. "Protein-bound uremic toxins in senescence and kidney fibrosis." *Biomedicines* 11 (2023): 2408.
42. Yang, Yi, Milos Mihajlovic, Manoe J. Janssen and Rosalinde Masereeuw. "The uremic toxin indoxyl sulfate accelerates senescence in kidney proximal tubule cells." *Toxins* 15 (2023): 242.
43. Glorieux, Griet, Sanjay K. Nigam, Raymond Vanholder and Francis Verbeke. "Role of the microbiome in gut-heart-kidney cross talk." *Circ Res* 132 (2023): 1064-1083.
44. Evenepoel, Pieter, Ruben Poesen and Björn Meijers. "The gut-kidney axis." *Pediatr Nephrol* 32 (2017): 2005-2014.
45. Vaziri, Nosratos D., Jun Yuan, Sohrab Nazetehrani and Zhenmin Ni, et al. "Chronic kidney disease causes disruption of gastric and small intestinal epithelial tight junction." *Am J Nephrol* 38 (2013): 99-103.
46. Mafra, Denise, Natália Borges, Livia Alvarenga and Marta Esgalhadó, et al. "Dietary components that may influence the disturbed gut microbiota in chronic kidney disease." *Nutrients* 11 (2019): 496.
47. Russell, Wendy R., Silvia W. Gratz, Sylvia H. Duncan and Grietje Holtrop, et al. "High-protein, reduced-carbohydrate weight-loss diets promote metabolite profiles likely to be detrimental to colonic health." *Am J Clin Nutr* 93 (2011): 1062-1072.
48. Ruiz-Ortega, Marta, Sandra Rayego-Mateos, Santiago Lamas and Alberto Ortiz, et al. "Targeting the progression of chronic kidney disease." *Nat Rev Nephrol* 16 (2020): 269-288.
49. Yarbeygi, Habib, Alexandra E. Butler, Stephen L. Atkin and Niki Katsiki, et al. "Sodium-glucose cotransporter 2 inhibitors and inflammation in chronic kidney disease: Possible molecular pathways." *J Cell Physiol* 234 (2019): 223-230.
50. Remuzzi, Giuseppe, Norberto Perico, Manuel Macia and Piero Ruggenenti. "The role of renin-angiotensin-aldosterone system in the progression of chronic kidney disease." *Kidney Int* 68 (2005): S57-S65.
51. Winiarska, Agata, Monika Knysak, Katarzyna Nabrdalik and Janusz Gumprecht, et al. "Inflammation and oxidative stress in diabetic kidney disease: The targets for SGLT2 inhibitors and GLP-1 receptor agonists." *Int J Mol Sci* 22 (2021): 10822.
52. Robledinos-Antón, Natalia, Raquel Fernández-Ginés, Gina Manda and Antonio Cuadrado. "Activators and inhibitors of NRF2: A review of their potential for clinical development." *Oxid Med Cell Longev* 2019 (2019).
53. "Exploring molecular targets in diabetic kidney disease." *Kidney Res Clin Pract* (2022).
54. Cantero-Navarro, Elena, Sandra Rayego-Mateos, Macarena Orejudo and Lucía Tejedor-Santamaria, et al. "Role of macrophages and related cytokines in kidney disease." *Front Med* 8 (2021): 688060.
55. Castillo-Rodríguez, Esmeralda, Raul Fernandez-Prado, Raquel Esteras and Maria Vanessa Perez-Gomez, et al. "Impact of altered intestinal microbiota on chronic kidney disease progression." *Toxins* 10 (2018): 300.
56. Wing, Maria R., Samir S. Patel, Ali Ramezani and Dominic S. Raj. "Gut microbiome in chronic kidney disease." *Exp Physiol* 101 (2016): 471-477.
57. Ramezani, Ali and Dominic S. Raj. "The gut microbiome, kidney disease and targeted interventions." *J Am Soc Nephrol* 25 (2014): 657.
58. Camerotto, Carla, Adamasco Cupisti, Claudia D'Alessandro and Fulvio Muzio, et al. "Dietary fiber and gut microbiota in renal diets." *Nutrients* 11 (2019): 2149.
59. Wiese, Gretchen N., Annabel Biruete, Ranjani N. Moorthi and Sharon M. Moe, et al. "Plant-based diets, the gut microbiota, and trimethylamine N-oxide production in chronic kidney disease: Therapeutic potential and methodological considerations." *J Ren Nutr* 31 (2021): 121-131.
60. Sumida, Keiichi, Wei Ling Lau, Csaba P. Kovesdy and Kamyar Kalantar-Zadeh, et al. "Microbiome modulation as a novel therapeutic approach in chronic kidney disease." *Curr Opin Nephrol Hypertens* 30 (2021): 75-84.
61. Gibson, Glenn R., Robert Hutkins, Mary Ellen Sanders and Susan L. Prescott, et al. "Expert consensus document: The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of prebiotics." *Nat Rev Gastroenterol Hepatol* 14 (2017): 491-502.
62. Hill, Colin, Francisco Guarner, Gregor Reid and Glenn R. Gibson, et al. "Activity of cecropin P1 and FA-LL-37 against urogenital microflora." *Nat Rev Gastroenterol Hepatol* 11 (2014): 506.

63. Swanson, Kelly S., Glenn R. Gibson, Robert Hutkins and Raylene A. Reimer, et al. "The International Scientific Association for Probiotics and Prebiotics (ISAPP) consensus statement on the definition and scope of synbiotics." *Nat Rev Gastroenterol Hepatol* 17 (2020): 687-701.
64. Cosola, Carmela, Maria Teresa Rocchetti, Ighli di Bari and Paola Maria Acquaviva, et al. "An innovative synbiotic formulation decreases free serum indoxyl sulfate, small intestine permeability and ameliorates gastrointestinal symptoms in a randomized pilot trial in stage IIIb-IV CKD patients." *Toxins* 13 (2021): 334.
65. Vaziri, Nosratola D. "Effect of synbiotic therapy on gut-derived uremic toxins and the intestinal microbiome in patients with CKD." *Clin J Am Soc Nephrol* 11 (2016): 199.
66. Liu, Sixiu, Hua Liu, Lei Chen and Shan-Shan Liang, et al. "Effect of probiotics on the intestinal microbiota of hemodialysis patients: A randomized trial." *Eur J Nutr* 59 (2020): 3755-3766.
67. Favero, Chiara, Laura Giordano, Silvia Maria Mihaila and Rosalinde Masereeuw, et al. "Postbiotics and kidney disease." *Toxins* 14 (2022): 623.
68. Mosca, Alexis, Ana Teresa Abreu Y Abreu, Kok Ann Gwee and Gianluca Ianiro, et al. "The clinical evidence for postbiotics as microbial therapeutics." *Gut Microbes* 14 (2022): 2117508.
69. Benech, Nicolas and Laetitia Koppe. "Is there a place for faecal microbiota transplantation in chronic kidney disease?." *Nephrol Dial Transplant* 37 (2022): 2303-2306.

How to cite this article: Anteyi, Emmanuel, Natarajan Ranganathan and Usha Vyas. "Targeting Inflammation to Slow CKD Progression- Role of Gut Microbiome Based Therapy." *J Nephrol Ther* 14 (2024): 481.